

Coronavirus Disease (SARS-CoV-2) and Neurocritical Care: Challenges for Neurosurgeons

Luis R Moscote-Salazar¹, Amit Agrawal², Moshir Rahman³

ABSTRACT

Neurocritical care is an important and integral part of neurosurgical management. In this global war without any smoke, it is a pivotal need for a holistic effort, as a health unit, in combating the virus. In this new era of SARS-CoV-2, neurosurgeons must acknowledge the insights regarding neurocritical care to safeguard our patients and quest for their best possible clinical outcome, working in harmony and sync with other clinical specialties. SARS-CoV-2 has very similar properties to the already known SARS-CoV, demonstrating in animal models its ability to invade neural tissues, with a high preponderance to neurons in the respiratory centers. The exact cause must be differentiated from nonspecific causes from those caused valid or indirect way by the infection, including infectious, para-infectious, and post-infectious encephalitis, hyper-coagulable states prompting stroke. To improve the quality of life of patients, roping in of the neurological manpower is mandatory in ICU, ward, research as well as neurological, neuropsychological, and neurocognitive rehabilitation including domiciliary care in search of a SARS-CoV-2 free world.

Keywords: Neurocritical care, Trauma, Trauma surgery.

Panamerican Journal of Trauma, Critical Care & Emergency Surgery (2021): 10.5005/jp-journals-10030-1317

Dear Editor,

Neurocritical care is an important and integral part of neurosurgical management. In this global war without any smoke, it is a pivotal need for a holistic effort, as a health unit, in combating the virus. In this new era of SARS-CoV-2, neurosurgeons must acknowledge the insights regarding neurocritical care to safeguard our patients and quest for their best possible clinical outcome, working in harmony and sync with other clinical specialties. Up to 20% of the patients have been seen to evolve to severe stages of the infection requiring intensive care support and management.¹ SARS-CoV-2 has very similar properties to the already known SARS-CoV, demonstrating in animal models its ability to invade neural tissues, with a high preponderance to neurons in the respiratory centers within the brain stem.² Exact cause must be differentiated from nonspecific causes (e.g., hypoxic encephalopathy and critical care neuropathy) from those caused valid or indirect way by the infection, including infectious, para-infectious, and post-infectious encephalitis, hyper-coagulable states prompting stroke.³ With the growing evidence of the neuroinvasive potential of this virus, it is indeed established that the virus has CSF penetration, and is attached to the SARS-CoV-2 receptor ACE2 in the brain, where it acts as a cell surface peptidase present on the surface of endothelial cells and neurons.⁴ Cerebral venous sinus thrombosis has got the fatal outcome. COVID patients presented with headaches with raised D-dimer, FDP, fibrinogen, and that is a strong predictor of cerebral thrombosis.⁵ Understanding the pathophysiology of COVID-19 and its impact on treating the neurological function in critical patients, like thromboprophylaxis, high-flow oxygen/extracorporeal membrane oxygenation, airway management, infection control, etc., are some of those.⁶ Age, sex, lymphocyte count, D-dimer, procalcitonin, brain natriuretic peptide, and respiratory support on SpO₂/FiO₂ were the prognosticators for critical COVID-19 patients.^{7,8}

European Academy of Neurology (EAN) scientific panel neurocritical care, stressed four challenges of neurocritical care during SARS-CoV-2 pandemic viz. (A) Neuro-invasiveness and involvement of the respiratory centers at the lower brainstem including pons is linked in respiratory failure implicating tapping

¹Department of Neurosurgery, Center for Biomedical Research, Faculty of Medicine, Universidad de Cartagena, Cartagena, Colombia

²Department of Neurosurgery, Narayana Medical College and Hospital, Nellore, Andhra Pradesh, India

³Neurosurgery Department, Holy Family Red Crescent Medical College Hospital, Dhaka, Bangladesh

Corresponding Author: MD Moshir Rahman, Neurosurgery Department, Holy Family Red Crescent Medical College Hospital, Dhaka, Bangladesh, Phone: +01880863944, e-mail: dr.tutul@yahoo.com

How to cite this article: Moscote-Salazar LR, Agrawal A, Moshir Rahman MD. Coronavirus Disease (SARS-CoV-2) and Neurocritical Care: Challenges for Neurosurgeons. *Panam J Trauma Crit Care Emerg Surg* 2021;10(2):89–90.

Source of support: Nil

Conflict of interest: None

cerebrospinal fluid obligatory to assay SARS-CoV-2 RNA; (B) Role of neurocritical care came into the picture as most patients are kept in a prone position with intermittent or continuous administration of muscle relaxants during multi-organ involvement with neuro-co-morbidities like a seizure, encephalitis, and cerebrovascular events (ischemic stroke or intracerebral hemorrhage), multifactorial encephalopathy, neuropathy, and myopathy; also post ICU-care syndrome with cognitive impairment, physical and psychiatric disabilities are reported. (C) Neurocritical care of traumatic brain injury, ischemic and hemorrhagic stroke, status epilepticus, neuro-immunological diseases should get equivalence, keeping in mind emergent issues of neuro-invasiveness of this virus; critical approach should get priority in co-incidence and finding causality. (D) SARS-CoV-2 ICU survivors should be on long-term follow-up and neurorehabilitation in the presence of neurological complications, post-ICU syndrome; subacute neurological complications, including Guillain-Barre syndrome, necrotizing encephalitis, etc., need evaluation. Thus, to improve the quality of life of ICU survivors,

roping in of the neurological manpower is mandatory in ICU, ward, research as well as neurological, neuropsychological, and neurocognitive rehabilitation including domiciliary care in search of a SARS-CoV-2 free world.⁹

REFERENCES

1. WHO, Report of the WHO-China Joint Mission on Coronavirus Disease 2019 (COVID-19) 2020 [Available from: [https://www.who.int/publications-detail/report-of-the-who-china-joint-mission-on-coronavirus-disease-2019-\(covid-19\)](https://www.who.int/publications-detail/report-of-the-who-china-joint-mission-on-coronavirus-disease-2019-(covid-19))].
2. Li YC, Bai WZ, Hashikawa T. The neuroinvasive potential of SARS-CoV2 may play a role in the respiratory failure of COVID-19 patients. *J Med Virol* 2020;92(6):552–555. DOI: 10.1002/jmv.25728.
3. Ellul MA, Benjamin L, Singh B, et al. Neurological associations of COVID-19. *Lancet Neurol* 2020;19(9):767–783. DOI: 10.1016/S1474-4422(20)30221-0.
4. Richardson PJ, Ottaviani S, Prell A, et al. CNS penetration of potential anti-COVID-19 drugs. *J Neurol*. 2020;267(7):1880–1882. DOI: 10.1007/s00415-020-09866-5.
5. Sasidharan PK. Cerebral vein thrombosis misdiagnosed and mismanaged. 2012;2012:210676. DOI: 10.1155/2012/210676.
6. Vanamoorthy P, Singh GP, Bidkar PU, et al. The Neurocritical Care Society of India (NCSI) and the Indian Society of Neuroanaesthesiology and Critical Care (ISNACC) joint position statement and advisory on the practice of neurocritical care during the COVID-19 pandemic. *J Neuroanaesthesiol Crit Care* 2020;7(03). DOI: 10.1055/s-0040-1714648.
7. Lu X, Jiang L, Chen T, et al. Continuously available ratio of SpO₂/FiO₂ serves as a noninvasive prognostic marker for intensive care patients with COVID-19. *Respir Res* 2020;21(1):194. DOI: 10.1186/s12931-020-01455-4.
8. Gavin W, Campbell E, Zaidi SA, et al. Clinical characteristics, outcomes and prognosticators in adult patients hospitalized with COVID-19. *Am J Infect Control* 2020;49(2):158–165. DOI: 10.1016/j.ajic.2020.07.005.
9. Helbok R, Öztürk S, COVID-19 Statement of the EAN Scientific Panel Neurocritical care 2020 [updated 2020/04/16/T13:57:35+00:00. Available from: <https://www.eanpages.org/2020/04/16/covid-19-statement-of-the-ean-scientific-panel-neurocritical-care/>.